

Claims:

- sub B17
1. A process for the preparation of ^aprotein hydrolysate from ^{defatted}soy flour using ^afungal protease, said process comprising: preparing ^{an}aqueous slurry of ^{said}defatted soy flour having 6-12% w/v of solid content; hydrolyzing the said slurry using fungal protease at pH 7-8 and temperature $43 \pm 5^\circ\text{C}$ for 1 to 3 hours to get 20-40% degree of hydrolysis (DH); further hydrolyzing using papain at temperature $53 \pm 5^\circ\text{C}$ for 0.5 to 1.5 hours under stirring till 30 - 45% DH is obtained; inactivating residual enzyme in a known manner; separating the solids and drying the clarified supernatant thus obtained to get protein hydrolysate.
 2. A process as claimed in claim 1, wherein the solid content in the slurry ranges from 8 - 12% w/v.
 3. A process as claimed in claim 1, wherein the fungal protease is obtained from *Aspergillus* sp.
 4. A process as claimed in claim 1, wherein *Aspergillus* is selected from the group comprising of *A. flavus*, *A. japonicus*, *A. niger* and *A. awamori*.
 5. A process as claimed in claim 1, wherein the protein hydrolysate is obtained by double enzyme hydrolysis.
 6. A process as claimed in claim 1, wherein the protein hydrolysate is obtained by hydrolyzing the slurry with proteolytic enzyme.
 7. A process as claimed in claim 1, wherein the fungal protease ranges from 0.4 to 0.5% w/w of the soy flour.
 8. A process as claimed in claim 1, wherein the protease hydrolysis is carried out at a pH of 7.2 to 7.6.

09811766-031901

- 09811766-031901
9. A process as claimed in claim 1, wherein the amount of papain ranges from 0.4 to 0.5% w/w of the soy flour.
 10. A process as claimed in claim 1, wherein the hydrolysate produced has decreased bitterness.
 11. A process as claimed in claim 1, wherein the threshold perception of bitterness is greater than 2g %.
 12. A process as claimed in claim 1, wherein the protein hydrolysate produced has low mineral content.
 13. A process as claimed in claim 1, wherein protein hydrolysate with 35 to 45% degree of hydrolysis (DH) is obtained in high yield from the raw material taken.
 14. A process as claimed in claim 1, wherein protein hydrolysate obtained has creamy color and a yield of 60-67.0% (on protein basis).
 15. A process as claimed in claim 1, wherein protein hydrolysate has 9.4% moisture, 10.5-11.0% nitrogen and 35-45% degree of hydrolysis.
 16. A process as claimed in claim 1, wherein the protein hydrolysate obtained has 20-23 trypsin inhibitor units/mg activity, 95 to 98% Nitrogen Solubility Index, 0.6 to 1.0% of salt content and 2 to 2.2 % bitterness recognition threshold.
 17. A process as claimed in claim 1, wherein lipoxxygenase and urease activities of the protein hydrolysate were not detectable.
 18. A process as claimed in claim 1, wherein the amino acid composition of the protein hydrolysate was similar to the amino acid makeup of starting material.

19. A protein hydrolysate obtained from soy flour, comprising 20 to 23 trypsin inhibitor units/mg activity, 95 to 98% Nitrogen Solubility Index, 0.6 to 1.0% salt content, 2 to 2.2% bitterness recognition threshold, 9.4% moisture, 10.5 to 11% nitrogen, and 35 to 45 % degree of hydrolysis.
20. A protein hydrolysate as claimed in claim 19, comprising undetectable lipoxigenase and urease activities.
21. A protein hydrolysate as claimed in claim 19, wherein the protein hydrolysate has similar acid makeup as that of the starting material.
22. A protein hydrolysate as claimed in claim 19, has cream color.

09811766-031901